

Abatacept (Orencia®) Prescriber Order and Referral Form

Pharmacists with the Froedtert Pharmacy will be supporting this patient's infusion medication management.

Fax completed form, insurance information and clinical documentation to 414-260-7368.

Patient Name: _____ Date of Birth: _____

Address: _____

Phone: _____ Height: _____ inches cm Weight: _____ lbs kg

Clinical Information

Primary Diagnosis Description: _____ ICD-10 Code(s): _____

Is this the first dose? Yes - Date of first dose _____ No - Date of next dose _____

Hepatitis B Status: Titer Date: _____ Positive Negative

TB Status PPD (negative) – date: _____ Active TB Unknown
 Last chest x-ray – date: _____ Other _____
 Past positive TB infection, course taken: _____

Abatacept (Orencia®) Prescription

Abatacept (Orencia®) refill as directed x 1 year

Initial Dose: Infuse 500 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight < 60 kg)
 Infuse 750 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight 60 to 100)
 Infuse 1000 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight > 100 kg)
 Other: _____

Maintenance Dose: Infuse 500 mg IV over 30 minutes every 4 weeks (patient weight < 60 kg)
 Infuse 750 mg IV over 30 minutes every 4 weeks (patient weight 60 to 100)
 Infuse 1000 mg IV over 30 minutes every 4 weeks (patient weight > 100 kg)
 Other: _____

Ancillary Orders

Anaphylaxis Orders

Treat per Froedtert Home and Specialty Infusion protocol.

Medication Orders

- Acetaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for fever or mild discomfort. Patient may decline.
- Diphenhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. Patient may decline.
- Methylprednisolone 40 mg IV push 20 minutes prior to infusion.
- Other: _____

IV Access and Flush Orders: RN to access and monitor venous infusion line as indicated below:

- Peripheral: Flush with 0.9% NS - 10 ml pre/post-use.
- Implanted Port or PICC: Flush with 0.9% NS - 10 ml pre/20 ml post-use and 10 ml pre/20 ml post-lab draw.

Lab Orders

- No labs ordered at this time.
- Other: _____

Skilled nurse to assess and administer and/or teach self-administration, where appropriate, via access device as indicated above. Nurse will provide ongoing support as needed. Refill above ancillary orders as directed x 1 year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature _____ Date _____ Time _____

Prescriber Name _____ Phone _____ Fax _____

Address _____ NPI _____

City, State, Zip _____ Office Contact _____



Original - Medical Records

Froedtert Home and Specialty Infusion
N86 W12999 Nightingale Way
Menomonee Falls, WI 53051
262-532-5040